

### **31. ZADATAK**

Koeficijent aktivnosti komponente(1) u dvokomponentnoj otopini ovisi, pri stalnom tlaku i temperaturi, o sastavu prema sljedećem empirijskom izrazu:

$$\ln \gamma_1 = ax_2^2 + bx_2^3 + cx_2^4 .$$

Parametri  $a$ ,  $b$  i  $c$  pritom ne ovise o sastavu.

Treba prirediti izraze koji opisuju ovisnost koeficijenta aktivnosti komponente(2), odnosno eksces Gibbsove energije o sastavu u istoj dvokomponentnoj otopini.

## GIBBS-DUHEMOVA JEDNADŽBA



Daje međuovisnost parcijalnih molarnih veličina u višekomponentnim sustavima:

$$\sum n_i d\bar{y}_i = 0$$

$$\sum x_i d\bar{y}_i = 0$$

Veza eksces Gibbsove energije i koeficijenta aktivnosti:

$$\frac{g^{\text{ex}}}{RT} = \sum x_i \ln \gamma_i$$



Logaritam koeficijenta aktivnosti jest parcijalna molarna veličina:

$$\sum x_i d \ln \gamma_i = 0$$

$$x_1 d \ln \gamma_1 + x_2 d \ln \gamma_2 = 0$$

$$x_2 d \ln \gamma_2 = -x_1 d \ln \gamma_1$$

$$d \ln \gamma_2 = -\frac{x_1}{x_2} d \ln \gamma_1$$

$$\int_{\ln \gamma_2(x_2=1)}^{\ln \gamma_2(x_2)} d \ln \gamma_2 = -\int_{x_2=1}^{x_2} \frac{x_1}{x_2} d \ln \gamma_1$$

$$x_1 = 1 - x_2$$

$$\frac{\partial \ln \gamma_1}{\partial x_2} = \frac{1}{\partial x_2} \partial (ax_2^2 + bx_2^3 + cx_2^4)$$

$$\frac{\partial \ln \gamma_1}{\partial x_2} = 2ax_2 + 3bx_2^2 + 4cx_2^3$$

$$d \ln \gamma_1 = (2ax_2 + 3bx_2^2 + 4cx_2^3) dx_2$$

Uvrštavanje:

$$\int_{\ln \gamma_2(x_2=1)}^{\ln \gamma_2(x_2)} d \ln \gamma_2 = -\int_{x_2=1}^{x_2} \frac{1-x_2}{x_2} (2ax_2 + 3bx_2^2 + 4cx_2^3) dx_2$$

Integriranje:

$$\ln \gamma_2 - 0 = -\int_{x_2=1}^{x_2} (1-x_2)(2a + 3bx_2 + 4cx_2^2) dx_2$$

$$\ln \gamma_2 - 0 = -\int_{x_2=1}^{x_2} [2a + (3b-2a)x_2 + (4c-3b)x_2^2 - 4cx_2^3] dx_2$$

$$\ln \gamma_2 = -\left(2ax_2 + \frac{3b-2a}{2}x_2^2 + \frac{4c-3b}{3}x_2^3 - \frac{4c}{4}x_2^4\right) \Big|_1^{x_2}$$

$$\ln \gamma_2 = -\left(2a(x_2 - 1) + \frac{3b - 2a}{2}(x_2^2 - 1) + \frac{4c - 3b}{3}(x_2^3 - 1) - \frac{4c}{4}(x_2^4 - 1)\right)$$

$$x_2 = 1 - x_1$$

$$\ln \gamma_2 = \left(a + \frac{3b}{2} + 2c\right)x_1^2 - \left(b + \frac{8c}{3}\right)x_1^3 + cx_1^4$$

Eksces Gibbsova energija:

$$\frac{g^{\text{ex}}}{RT} = x_1 \ln \gamma_1 + x_2 \ln \gamma_2$$

$$\frac{g^{\text{ex}}}{RT} = x_1(ax_2^2 + bx_2^3 + cx_2^4) + x_2 \left[ \left(a + \frac{3b}{2} + 2c\right)x_1^2 - \left(b + \frac{8c}{3}\right)x_1^3 + cx_1^4 \right]$$

$$\frac{g^{\text{ex}}}{RT} = x_1 x_2 \left[ ax_2 + bx_2^2 + cx_2^3 + \left(a + \frac{3b}{2} + 2c\right)x_1 - \left(b + \frac{8c}{3}\right)x_1^2 + cx_1^3 \right]$$

$$\frac{g^{\text{ex}}}{RT} = x_1 x_2 \left[ a + b + c - \left(\frac{b}{2} + c\right)x_1 + \frac{c}{3}x_1^2 \right]$$

Korektan izraz za eksces Gibbsovu energiju!